



Measles, mumps and rubella vs diphtheria-tetanus-acellular-pertussis-inactivated-polio-Haemophilus influenzae type b as the most recent vaccine and risk of early 'childhood asthma'

Rieckmann, Andreas; Hærskjold, Ann; Benn, Christine Stabell; Aaby, Peter; Lange, Theis; Sørup, Signe

Published in:
International Journal of Epidemiology

DOI:
[10.1093/ije/dyz062](https://doi.org/10.1093/ije/dyz062)

Publication date:
2019

Document version
Peer reviewed version

Citation for published version (APA):
Rieckmann, A., Hærskjold, A., Benn, C. S., Aaby, P., Lange, T., & Sørup, S. (2019). Measles, mumps and rubella vs diphtheria-tetanus-acellular-pertussis-inactivated-polio-Haemophilus influenzae type b as the most recent vaccine and risk of early 'childhood asthma'. *International Journal of Epidemiology*, 48(6), 2026-2038. <https://doi.org/10.1093/ije/dyz062>

Measles, mumps, and rubella vs diphtheria-tetanus-acellular-pertussis-inactivated-polio-*Haemophilus-influenzae*-type-b as the most recent vaccine and risk of early “childhood asthma”

Authors: Andreas Rieckmann^{1,2}, Ann Hærskjold³, Christine Stabell Benn^{1,4}, Peter Aaby⁵, Theis Lange^{6,7}, and Signe Sørup^{1,8}.

Affiliations: ¹Research Center for Vitamins and Vaccines (CVIVA), Bandim Health Project, Statens Serum Institut, Copenhagen, Denmark, ² Section of Epidemiology, Department of Public Health, The University of Copenhagen, Copenhagen, Denmark ³Department of Dermatology, Bispebjerg Hospital, Copenhagen, Denmark, ⁴OPEN, Odense University Hospital/Institute of Clinical Research, University of Southern Denmark, Odense, Denmark, ⁵Bandim Health Project, Indepth Network, Bissau, Guinea-Bissau, ⁶Section of Biostatistics, Department of Public Health, University of Copenhagen, Copenhagen, Denmark, ⁷Center for Statistical Science, Peking University, China. ⁸Department of Clinical Epidemiology, Aarhus University, Aarhus N, Denmark.

Key words: Asthma; Immunity, Heterologous; Immunization; Measles-Mumps-Rubella Vaccine; Non-specific effects of vaccines; Infant, Asthma; Vaccine-sequence; Trained innate immunity

Corresponding author: Andreas Rieckmann

Conflicts of interest: none

Journal: International Journal of Epidemiology

Abstract: 240 words, Manuscript: 3046 words, Figures: 3, Tables: 4, References: 38, Supplementary material: Supplementary figures: 7, Supplementary tables: 9, Supplementary information: 2.

Funding

This work was supported by the Danish National Research Foundation (DNRF), which supports the Research Center for Vitamins and Vaccines [DNRF108]. AR was supported by an unrestricted Faculty of Health Sciences-scholarship from University of Southern Denmark. SS holds a grant from the Danish Council for Independent Research [DFF – 4183-00316].

Acknowledgement

The authors thank professor Poul-Erik Lund Kofoed for his valuable comments on childhood asthma. We also want to thank the referees reviewing our paper for many constructive advises.

Abstract

Background and objective: Live vaccines may have beneficial non-specific effects. We tested whether the live measles, mumps and rubella (MMR) vaccine compared with the non-live diphtheria-tetanus-acellular-pertussis-inactivated-polio-*Haemophilus-influenzae*-type-b (DTaP-IPV-Hib) vaccine as the most recent vaccine was associated with less childhood asthma and fewer acute hospital contacts for childhood asthma among boys and girls.

Methods: This study is a nationwide register-based cohort study of 338 761 Danish children born 1999-2006. We compared 1) the incidence of first-registered childhood asthma based on hospital contacts and drug prescriptions and 2) the incidence of severe asthma defined as acute hospital contacts for childhood asthma between 15 and 48 month among children who latest had received 3 doses of DTaP-IPV-Hib and then MMR with children who latest had received 3 doses of DTaP-IPV-Hib.

Results: For boys, following the recommended vaccine schedule of MMR after DTaP-IPV-Hib3 compared with DTaP-IPV-Hib3 as the latest vaccine, MMR was associated with 8.0 (95% confidence interval 3.9-12.2) fewer childhood asthma cases per 1,000 boys, corresponding to 10% (5-15%) reduction in the cumulative incidence of childhood asthma. MMR, when given last, was also associated with 16.3 (12.7-19.9) fewer acute hospital admissions for childhood asthma per 1,000 boys, corresponding to a 27% (22-31%) reduction in the cumulative incidence. No associations were seen for girls.

Conclusion: MMR may have a protective effect against childhood asthma for boys. This calls for an understand of whether non-specific effects of vaccines can be used to optimise our vaccine programmes.

Key Messages

- The live measles, mumps and rubella (MMR) vaccine as the most recently received vaccine may reduce childhood asthma.
- In a Danish population of children aged 15-48 months, the measles, mumps and rubella vaccine compared with the non-live diphtheria-tetanus-acellular-pertussis-inactivated-polio-haemophilus-influenzae-type-b vaccine (DTaP-IPV-Hib) as the most recent vaccine was associated with less childhood asthma and less severe asthma for boys.
- Timely MMR after DTaP-IPV-Hib may protect one in 124 (95% CI 82 to 260) boys in the age between 15-48 months from childhood asthma.
- This potential calls for a better understanding of non-specific effects of vaccines.

1 Introduction

2 Vaccines may have beneficial non-specific effects (NSEs).(1-4) Live vaccines affect the innate
3 immune system beneficially leading to increased response to unrelated pathogens(2) and are
4 associated with lower morbidity and mortality from non-targeted diseases. (1,3-7) Studies
5 suggest that live vaccines induce a T-helper 1 cell dominant shift reducing atopic
6 diseases.(4,8,9) In contrast, non-live vaccines can induce tolerance of the innate immune system
7 leading to decreased responses to unrelated pathogens(2) and have been associated with
8 higher morbidity and mortality from non-targeted diseases.(1,3-7) The non-specific effects of
9 live vaccines are most pronounced as long as a given vaccine is the most recent vaccine and
10 may vary by sex.(4,8,9)

11 Studies of the live measles, mumps, and rubella (MMR) vaccine and childhood asthma have
12 reported preventive effects(10-13), detrimental effects(14) or no effect(15-19). However,
13 vaccination sequence was not assessed in any of the studies (10-19) and sex-differential effects
14 was only assessed in one study(11).

15 The main objective of this study was to test the hypothesis that the live MMR vaccine *as the*
16 *most recent vaccine* compared with the non-live diphtheria-tetanus-acellular pertussis-polio-
17 *Haemophilus Influenzae* type b (DTaP-IPV-Hib) vaccine is associated with a lower incidence of
18 childhood asthma defined by hospital contacts and drug prescriptions in Danish children aged
19 15 to 48 months (hypothesis 1). Our secondary objective was to test whether MMR was

associated with a lower incidence of *severe* childhood asthma defined by acute hospital contacts (hypothesis 2). We examined whether associations differed by sex.

Method

This study is a register-based cohort study of Danish children followed from the age of 15 months until the age of 48 months for the risk of childhood asthma according to whether their last vaccine received was DTaP-IPV-Hib. Linkage across Danish nation-wide registers is possible using Danish personal identification numbers assigned to all new-borns. We used information from The Danish Civil Registration System, the Danish National Health Service Register, the Danish National Patient Register, the Danish National Prescription Register, the Danish Medical Birth Register, and information from Statistics Denmark (Supplementary information 1).

Study base and vaccinations

To reduce interference from other vaccines than MMR and DTaP-IPV-Hib, the study base was children born in Denmark from 1 July 1999 until and including 30 September 2006 and followed from 15 months of age until 48 months of age (Supplementary figure 1). For this study base, the Danish childhood vaccination programme until 48 months of age was three doses of the non-live DTaP-IPV-Hib at 3, 5, and 12 months of age and the first dose of the live MMR at 15 months of age.

37 In Denmark, vaccines are administered free-of-charge by the general practitioners. For the
38 purpose of reimbursement, information about vaccine type, dose, and week of vaccination is
39 collected in the Danish National Health Service Register.(20)

40 **Childhood asthma**

41 Asthma in young children may be differentiated by multiple phenotypes(21) and multiple ways
42 of phenotyping.(22) Asthma, asthmatic bronchitis and infection induced wheezing in young
43 children are difficult to separate using register data, hence we used the term “childhood
44 asthma” to cover the aforementioned diseases.

45 Childhood asthma was defined by information from two sources: 1) the Danish National Patient
46 Register(23) holds information on hospital contacts coded with the 10th version of the
47 international classification of diseases (ICD-10); 2) The Danish National Prescription Register(24)
48 holds information about all redeemed prescriptions in Danish pharmacies.

49 The main outcome (hypothesis 1) was the incidence of first-registered childhood asthma
50 between 15 to 48 months of age defined as the first registration of a hospital contacts
51 (outpatients, inpatients, and emergency room) according to ICD-10 codes J45 (*Asthma*) and J46
52 (*Status asthmaticus*) or the first prescription of inhaled glucocorticoids (R03BA) or montelukast
53 (R03DC03) (at least 2 prescriptions of either glucocorticoids or montelukast within 12 months).
54 Our second outcome (hypothesis 2) was severe childhood asthma between 15 to 48 months of
55 age defined as the incidence of all acute hospital contacts (inpatients and emergency room

visits) with ICD-10 codes J45 or J46. Allowing for recurrent events, adjacent acute hospital contact periods within 5 days of last hospital discharge counted as one acute hospital contact.

Inclusion and exclusion criteria

To ensure that the study population consisted of children, who followed the vaccination schedule before the baseline of our study, only children who had received the first two doses of DTaP-IPV-Hib before the age of 11 months were included. For hypothesis 1, we excluded children who had been registered with childhood asthma before 15 months of age, while this restriction did not apply for hypothesis 2, since it focused on repeated events (Figure 1 and Supplementary figure 2).

Sequence cohorts

A proportion of Danish children deviate from the recommended schedule of the childhood vaccination programme. This variation allows us to investigate the robustness of the analyses in three partly overlapping vaccine sequence cohorts (Figure 1); *recommended sequence* (DTaP-IPV-Hib1-3-then-MMR vs DTaP-IPV-Hib1-3 only), *early MMR sequence* (DTaP-IPV-Hib1-2-then-MMR vs DTaP-IPV-Hib1-2 only) and *interrupted sequence* (DTaP-IPV-Hib1-2-then-MMR vs DTaP-IPV-Hib1-2-then-MMR-then-DTaP-IPV-Hib3).

Follow-up

73 For hypothesis 1, all children entered the study at the first Sunday after they turned 15 months
74 of age or when their vaccination sequence qualified them for one of the sequence cohorts
75 described above, whichever occurred last. All children left the study: a) at the date of a
76 registered childhood asthma event, b) at 48 months of age, c) when receiving more than one
77 vaccine in one day or receiving unrelated vaccines, or d) 12 months before emigrating or dying,
78 to allow two prescriptions to be registered within a year, whichever occurred first. For
79 hypothesis 2, the same entry and exit criteria applied, except that children registered with an
80 acute hospital contact for childhood asthma left the study for 5 days and re-entered the study
81 to make sure that we did not count one episode several times.

82 **Statistical methods**

83 We suspected that hospital admissions due to infectious diseases could both affect vaccination
84 status(25,26) and later childhood asthma(27,28); furthermore hospital admissions due to
85 infectious diseases have been shown to be reduced by prior MMR vaccination(4,29) (illustrated
86 in Supplementary figure 3). This is known as time-varying confounding. We attempted to
87 control for this potential issue using weights, which we updated each week based on 22
88 covariates (Supplementary information 1 and 2). Such an approach is well-established and aims
89 at creating comparable comparison groups at any given time during follow up.(30,31) We have
90 added a technical description in Supplementary information 2.

We calculated the crude and weighted cumulative incidence rates and reported the cumulative incidence rate reduction per 1,000 children for each group at 48 months of age ($wIR_{reduction_{48}}$) and cumulative incidence rate ratios at 48 months of age. Standard errors were calculated using robust estimators and confidence intervals were estimated with 2.5% and 97.5% quantiles using 100,000 simulations. The analyses assumed independent observations. A spline function of the crude and weighted cumulative incidence rates were plotted for each analysis. We investigated whether the $wIR_{reduction_{48}}$ were different for girls and boys. Additionally, to compare with hazards ratios often reported in the medical literature, we used a weighted Cox proportional hazards model to estimate hazard ratios. The weighted Cox proportional hazards model had age as the underlying time scale and was stratified by week (Sunday to Saturday) of birth thereby adjusting for the effect of age and calendar time. We tested the assumption of proportional hazards using Schoenfeld residuals.

Sensitivity analysis

Since infections may affect childhood asthma(27,28) and MMR has been associated with fewer infections in the same cohort(4), we conducted a sensitivity analysis excluding hospital admissions with childhood asthma co-occurring with hospital admissions for an infectious disease, to ensure we did not partly reproduce earlier results.

If childhood asthma was registered using two prescriptions within one year, we used the first date of two prescriptions. Thus, the registered childhood asthma date was conditioned on a

future prescription, which could cause a collider bias introducing a non-causal association between MMR vaccination and the first prescription for childhood asthma. We conducted a sensitivity analysis using the second date of prescriptions.

Furthermore, we conducted a sensitivity analysis of the effect of DTaP-IPV-Hib 3 vs DTaP-IPV-Hib 2 from 12 to 15 months of age under the hypothesis that the incidence of new-onset childhood asthma and severe asthma would not differ. If DTaP-IPV-Hib 3 is associated with a reduced incidence of asthma compared with DTaP-IPV-Hib 2 in spite of adjustment for potential confounding, this would suggest that we have been unable to fully control for confounding i.e. that the most healthy children receive the next vaccine (residual confounding of a "healthy vaccinee effect" -bias) Data was analysed using Stata 14.0 and R 3.3.1 with version 1.0-11 for the ipw package. Register based studies do require ethical approval by the Danish Central Scientific Ethics Committee.

Results

From the study base of 467,919 children, 338,761 children were eligible for studying hypothesis 1 (Figure 1) and 364,270 children were eligible for hypothesis 2 (Supplementary figure 2). The recommended sequence included the majority of risk time with 773,054 person years of follow up (pyrs), the early MMR sequence had 50,918 pyrs, and the interrupted sequence had 36,788 pyrs.

128 Most shifts in vaccination sequence cohorts happened between 15-20 months of age
129 (Supplementary figure 4 and 5). Childhood asthma peaked at 8 months of age, and boys have a
130 higher incidence than girls (Supplementary figure 6). Among the recommended sequence for
131 hypothesis 1, 18 776 children were registered with childhood asthma during follow-up (of
132 which; 25% based on hospital contacts, 74% based on glucocorticoids, and 1% based on
133 montelukast).

134 Covariates were approximately similarly distributed between vaccination sequence groups at 16
135 months of age (Supplementary table 1). However, children who had received only DTaP-IPV-
136 Hib 1 + 2 at 16 months of age tended to have a skewed distribution compared with the
137 remaining vaccination sequence groups especially regarding single parenthood, family income,
138 maternal education, other children in the household, mother smoking during pregnancy, and
139 previous infectious diseases and prescribed antibiotics.

140 For hypothesis 1, MMR compared with DTaP-IPV-Hib as most recent vaccine was associated
141 with a weighted cumulative incidence reduction of childhood asthma per 1,000 children at 48
142 months of age ($wIR_{reduction_{48}}$) of 4.5 (95% confidence interval (CI) 1.7 to 7.2), 3.8 (95% CI -3.3
143 to 10.8), and 22.9 (95% CI 3.1 to 42.7) for *the recommended, early, and interrupted sequence*
144 *respectively* (Table 1). The incidence diverged mostly in the first part of follow-up (Figure 2).
145 The incidence difference tended to be larger among boys than girls (Table 2).

For hypothesis 2, MMR compared with DTaP-IPV-Hib as the last received vaccine was associated with a $wIR_{reduction_{48}}$ of 8.2 (95% CI 6.0 to 10.5), 13.4 (95% CI 7.3 to 19.4), and 1.7 (95% CI -11.8 to 15.2) for the rate of acute hospital contacts for childhood asthma in the *recommended, early and interrupted sequence* (Table 3 and figure 3). Again, the association was strongest among boys compared with girls in the recommended sequence (Table 4).

When analysing the hypotheses using hazard ratios, the assumption of proportional hazards was not met for hypothesis 1 (Supplementary table 2) but was more stable for hypothesis 2 (Supplementary table 3).

Excluding concurrent hospital contacts for infectious diseases and using the date of the second prescription provided similar estimates (Supplementary tables 4, 5 and 6).

The weighted cumulative incidence difference from 12 to 15 months of age per 1,000 children for DTaP-IPV-Hib 3 compared with DTaP-IPV-Hib 2 for the hypothesis 1 and 2 outcomes were respectively 2.7 (95% CI 1.6 to 3.8) and 0.6 (95% CI -0.1 to 1.3). The association tended to be stronger among boys than girls (Supplementary tables 7 and 8).

Discussion

Among boys but not girls, MMR vaccination compared with DTaP-IPV-Hib as the last received vaccine across cohorts was associated with a lower incidence of childhood asthma and fewer acute hospital contacts for childhood asthma.

164 **Strengths and weaknesses**

165 This was a large nationwide study and we applied state of the art statistical methods, the
166 weighted approach, to handle potential time-varying confounding.(32) The weights were not
167 extreme so no single individual affected the results disproportionately (Supplementary figure 7).
168 The 95% confidence interval for the incidence rate difference was calculated assuming
169 independent observations, which is a limitation as children have contributed risk time to both
170 vaccination groups.

171 By ensuring that children had followed the vaccination programme (received DTaP-IPV-Hib 1
172 and 2 before 11 months of age) we increased homogeneity of the study population(33) and by
173 including many potential confounders, we may to some extent have controlled for unmeasured
174 confounding though it cannot be ruled out. Potential time-varying confounding due to
175 children' s health status was modelled by previous chronic diseases as well as hospital
176 admissions and antibiotic use for infectious diseases. The comparison between DTaP-IPV-Hib 3
177 and DTaP-IPV-Hib 2 for the hypothesis 1 and to some degree hypotheses 2 suggests that there
178 may be confounding not fully adjusted for (residual confounding of a "*healthy vaccinee*
179 *effect*" -bias). However, in the *interrupted sequence* cohort, children who received the next
180 vaccine (DTaP-IPV-Hib 3 after MMR) had a higher risk for childhood asthma compared with
181 those who had only received MMR, which indicates that a "*healthy vaccinee effect*" -bias does
182 not fully explain our results Receiving the MMR vaccine as the last vaccine was associated with
183 approximately similar cumulative incidence rates across sequences, while DTaP-IPV-Hib in

184 contrast was associated with greater cumulative incidence rates. As this pattern was
185 independent of which sequence children followed, it suggests that both vaccines affect the risk
186 of childhood asthma respectively beneficially and detrimentally.

187 MMR vaccination coverage in Denmark may be underreported,(34) which would increase
188 proportionally with time if MMR vaccinated children stay categorised as DTaP-IPV-Hib 3
189 vaccinated thus biasing the estimate towards null over time – a pattern we also observed. The
190 inclusion criteria DTaP-IPV-Hib 1 and 2 before 11 months of age decreased the misclassification
191 of the reference groups in the *recommended sequence* and *early MMR sequence*.
192 Unsystematic underreporting would also cause conservative estimates. Simultaneous
193 administration of vaccines was outside the scope of our analysis.

194 Neither of our sensitivity analyses indicated that we should have reached other conclusions.
195 Further analyses indicated that it was the first prescription of childhood asthma prescription
196 rather than the criteria of two prescriptions within one year that captured the effect of MMR on
197 childhood asthma (analysis not shown).

198 **Studies on MMR and childhood asthma**

199 No randomised trial has studied the effect of MMR on the risk of childhood asthma; nine
200 diverse observational studies on the effect of MMR or measles vaccines on childhood asthma
201 show conflicting results (Supplementary table 9). (10-19) The nine studies vary from large
202 prospective cohort studies to small cross-sectional studies and represent children from 11

countries (two sets of studies partly overlap in study populations[(16,17) and (18,19)] and one study partly overlap with the study population of our study(11)). The studies were further heterogenic with regards to outcome definition, age group and sequence of vaccines. Specifically, our study indicated that the effect of vaccine status on the incidence of childhood asthma decreased with age, making comparisons for unlike age groups problematic. Our study is the first study to strictly investigate the effect of having MMR as the most recent vaccine on the risk of childhood asthma. Not differentiating the vaccine sequence mixes effects of multiple vaccine combinations,(35) a problem, which was highlighted in a recent WHO review of non-specific effects of vaccines(3) and also supported by this study. Furthermore, applying the inclusion criteria where all children had received the first two doses of DTaP-IPV-Hib before 11 months of age increased the exchangeability of the comparison groups in our study compared with the previous studies. Despite differences in design, the study overlapping with the study population of our study found similar effects of MMR on the use of glucocorticoids and for hospital admissions due to asthma.(11) However, they did not find similar sex differential effects for hospital admissions due to asthma, which may have been masked by the effects of other vaccines.

Biological mechanisms

Successful immunisation by live vaccines, like natural infections, has been shown to shift from a predominant and often hyper reactive T-helper 2 cell to being T-helper 1 cell predominant that

222 allows an accelerated clearance of infections,(1) and thus a potentially reduced risk of
223 childhood asthma.

224 MMR compared with DTaP-IPV-Hib as the most recent vaccine has been associated with fewer
225 admissions for infectious diseases.(4) Since persistent respiratory infections may destroy
226 epithelial cells of the respiratory tract making children with childhood asthma more prone to
227 severe episodes(36) and as respiratory infections can function as irritants exaggerating wheeze
228 attacks(9), then a reduction in infectious diseases could lead to a lower risk of childhood
229 asthma.

230 Recent cutting edge immunological research show that live vaccines can train the innate
231 immune system for swifter clearance of non-related pathogens possible through epigenetic
232 reprogramming.(2) In accordance with our results indicating that males benefited more from
233 the MMR vaccine, a new immunological study of Gambian children showed that measles
234 vaccine was associated with enhanced proinflammatory innate responses for males but not for
235 females possibly by modifying signaling via Toll-like receptor 4 (TLR4).(37) The duration of
236 trained immunity is yet unknown, but according to our data the absolute risk difference was
237 proportionally largest the first 10-15 months after vaccination and waned off (Figures 2 and 3).

238 **Conclusion**

239 Our study indicated that boys experience less childhood asthma and fewer acute hospital
240 contacts for childhood asthma when the live MMR vaccine compared with the non-live DTaP-

241 IPV-Hib vaccine was the most recent vaccine. Data did not convincingly suggest an effect for
242 girls. For now, these results are only indications, thus we hope other research groups will test
243 our findings that MMR compared with DTaP-IPV-Hib as most recent vaccine reduces the risk of
244 childhood asthma among boys and investigate why girls may not experience the same benefit.
245 If supported, the question of whether a beneficial effect of MMR would persist or be countered
246 by a subsequent non-live vaccine should be addressed.

247 If our estimates are correct then for 1,000 boys following the recommended vaccine schedule
248 and receiving MMR timely after DTaP-IPV-Hib, MMR prevents 8.0 (95% CI 3.9 to 12.2)
249 childhood asthma cases between 15 months and 48 months. Thus, one childhood asthma case
250 is prevented for each 124 (95% CI 82 to 260) MMR vaccinations administered. Though most
251 childhood asthma cases occurred before 15 months of age in this study, potential immune
252 modulation via MMR or other approaches for preventing childhood asthma calls for a better
253 understanding of non-specific effects of vaccines.(38)

References

1. Benn CS, Netea MG, Selin LK, Aaby P. A small jab - a big effect: nonspecific immunomodulation by vaccines. *Trends Immunol.* 2013 Sep;34(9):431–9.
2. Netea MG, Joosten LAB, Latz E, Mills KHG, Natoli G, Stunnenberg HG, et al. Trained immunity: A program of innate immune memory in health and disease. *Science.* 2016 Apr 22;352(6284):aaf1098.
3. Higgins JPT, Soares-Weiser K, López-López JA, Kakourou A, Chaplin K, Christensen H, et al. Association of BCG, DTP, and measles containing vaccines with childhood mortality: systematic review. *BMJ.* 2016 Oct 13;355:i5170.
4. Sorup S, Benn CS, Poulsen A, Krause TG, Aaby P, Ravn H. Live vaccine against measles, mumps, and rubella and the risk of hospital admissions for nontargeted infections. *JAMA.* 2014 Feb 26;311(8):826–35.
5. Aaby P, Ravn H, Fisker AB, Rodrigues A, Benn CS. Is diphtheria-tetanus-pertussis (DTP) associated with increased female mortality? A meta-analysis testing the hypotheses of sex-differential non-specific effects of DTP vaccine. *Trans R Soc Trop Med Hyg.* 2016 Dec;110(10):570–81.
6. Mogensen SW, Andersen A, Rodrigues A, Benn CS, Aaby P. The Introduction of Diphtheria-Tetanus-Pertussis and Oral Polio Vaccine Among Young Infants in an Urban African Community: A Natural Experiment. *EBioMedicine.* 2017 Mar;17:192–8.
7. Bardenheier BH, McNeil MM, Wodi AP, McNicholl JM, DeStefano F. Risk of Nontargeted Infectious Disease Hospitalizations Among US Children Following Inactivated and Live Vaccines, 2005–2014. *Clin Infect Dis.* 2017 May 6.
8. Aaby P, Jensen H, Samb B, Cisse B, Sodemann M, Jakobsen M, et al. Differences in female-male mortality after high-titre measles vaccine and association with subsequent vaccination with diphtheria-tetanus-pertussis and inactivated poliovirus: reanalysis of West African studies. *Lancet.* 2003 Jun 28;361(9376):2183–8.
9. Douwes J, Pearce N. Asthma and the westernization 'package'. *Int J Epidemiol.* 2002 Dec;31(6):1098–102.
10. Jedrychowski W, Maugeri U, Jedrychowska-Bianchi I. Prospective epidemiologic study on respiratory diseases in children and immunization against measles. *Int J Occup Med Environ Health.* 2004;17(2):255–61.

11. Hviid A, Melbye M. Measles-mumps-rubella vaccination and asthma-like disease in early childhood. *Am J Epidemiol*. 2008 Dec 1;168(11):1277–83.
12. Grabenhenrich LB, Gough H, Reich A, Eckers N, Zepp F, Nitsche O, et al. Early-life determinants of asthma from birth to age 20 years: a German birth cohort study. *J Allergy Clin Immunol*. 2014 Apr;133(4):979–88.
13. Timmermann CAG, Osuna CE, Steuerwald U, Weihe P, Poulsen LK, Grandjean P. Asthma and allergy in children with and without prior measles, mumps, and rubella vaccination. *Pediatr Allergy Immunol*. 2015 Dec;26(8):742–9.
14. McKeever TM, Lewis SA, Smith C, Hubbard R. Vaccination and allergic disease: a birth cohort study. *Am J Public Health*. 2004 Jun;94(6):985–9.
15. Wickens K, Crane J, Kemp T, Lewis J, Kennedy J, D'Souza W, et al. A case-control study of risk factors for asthma in New Zealand children. *Aust N Z J Public Health*. 2001;25(1):44–9.
16. Maher JE, Mullooly JP, Drew L, DeStefano F. Infant vaccinations and childhood asthma among full-term infants. *Pharmacoepidemiol Drug Saf*. 2004 Jan;13(1):1–9.
17. DeStefano F, Gu D, Kramarz P, Truman BI, Iademarco MF, Mullooly JP, et al. Childhood vaccinations and risk of asthma. *Pediatr Infect Dis J*. 2002 Jun;21(6):498–504.
18. Flöistrup H, Flöistrup H, Swartz J, Swartz J, Bergström A, Bergström A, et al. Allergic disease and sensitization in Steiner school children. *J Allergy Clin Immunol*. 2006 Jan;117(1):59–66.
19. Rosenlund H, Bergström A, Alm JS, Swartz J, Scheynius A, van Hage M, et al. Allergic disease and atopic sensitization in children in relation to measles vaccination and measles infection. *Pediatrics*. 2009 Mar;123(3):771–8.
20. Andersen JS, Olivarius NDF, Krasnik A. The Danish National Health Service Register. *Scand J Public Health*. 2011 Jul;39(7 Suppl):34–7.
21. Stein RT, Stein RT, Holberg CJ, Holberg CJ, Morgan WJ, Morgan WJ, et al. Peak flow variability, methacholine responsiveness and atopy as markers for detecting different wheezing phenotypes in childhood. *Thorax*. 1997 Nov;52(11):946–52.
22. Spycher BD, Silverman M, Kuehni CE. Phenotypes of childhood asthma: are they real? *Clin Exp Allergy*. 2010 Aug;40(8):1130–41.
23. Schmidt M, Schmidt SAJ, Sandegaard JL, Ehrenstein V, Pedersen L, Sørensen HT. The Danish National Patient Registry: a review of content, data quality, and research potential. *CLEP*.

2015;7:449–90.

24. Wallach Kildemoes H, Toft Sorensen H, Hallas J. The Danish National Prescription Registry. *Scand J Public Health*. 2011 Jul 20;39(7 Suppl):38–41.
25. Virtanen M, Peltola H, Paunio M, Heinonen OP. Day-to-day reactogenicity and the healthy vaccinee effect of measles-mumps-rubella vaccination. *Pediatrics*. 2000 Nov;106(5):E62.
26. Fine PEM, Chen RT. Confounding in Studies of Adverse Reactions to Vaccines. *Am J Epidemiol*. 1992 Aug 1;121–35.
27. Sigurs N, Aljassim F, Kjellman B, Robinson PD, Sigurbergsson F, Bjarnason R, et al. Asthma and allergy patterns over 18 years after severe RSV bronchiolitis in the first year of life. *Thorax*. 2010 Dec;65(12):1045–52.
28. Johnston SL, Pattemore PK, Sanderson G, Smith S, Lampe F, Josephs L, et al. Community study of role of viral infections in exacerbations of asthma in 9-11 year old children. *BMJ*. 1995 May 13;310(6989):1225–9.
29. Sorup S, Jensen AKG, Aaby P, Benn CS. Revaccination with Measles-mumps-rubella Vaccine and Infectious Disease Morbidity: A Danish Register-based Cohort Study. *Clin Infect Dis*. 2018 May 26.
30. Robins JM, Hernan MA, Brumback B. Marginal structural models and causal inference in epidemiology. *Epidemiology*. 2000 Sep;11(5):550–60.
31. Hernan MA, Brumback B, Robins JM. Marginal structural models to estimate the causal effect of zidovudine on the survival of HIV-positive men. *Epidemiology*. 2000 Sep;11(5):561–70.
32. Hernan MA, Robins JM. *Causal Inference*. Boca Raton: Chapman & Hall/CRC, forthcoming; 2017. 45 p.
33. Pizzi C, De Stavola BL, Pearce N, Lazzarato F, Ghiotti P, Merletti F, et al. Selection bias and patterns of confounding in cohort studies: the case of the NINFEA web-based birth cohort. *J Epidemiol Community Health*. 2012 Nov;66(11):976–81.
34. Holt N, Mygind A, Bro F. Danish MMR vaccination coverage is considerably higher than reported. *Dan Med J*. 2017 Feb;64(2).
35. Benn CS, Sorup S. Commentary: BCG has no beneficial non-specific effects on Greenland. An answer to the wrong question? *Int J Epidemiol*. 2016 Dec 1;45(6):2131–3.
36. Martinez FD. Role of respiratory infection in onset of asthma and chronic obstructive

pulmonary disease. Clin Exp Allergy. 1999 Jun;29 Suppl 2:53–8.

37. Noho-Konteh F, Adetifa JU, Cox M, Hossin S, Reynolds J, Le MT, et al. Sex-Differential Non-Vaccine-Specific Immunological Effects of Diphtheria-Tetanus-Pertussis and Measles Vaccination. Clin Infect Dis. 2016 Nov 1;63(9):1213–26.
38. Pollard AJ, Finn A, Curtis N. Non-specific effects of vaccines: plausible and potentially important, but implications uncertain. Arch Dis Child. 2017 May 13.